

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 647 712 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 93116221.8

(51) Int. Cl.⁶: **C12N 15/54, C12P 19/40,
C12N 15/70, C12N 1/21,
/(C12N1/21,C12R1:19)**

(22) Date of filing: 07.10.93

(43) Date of publication of application:
12.04.95 Bulletin 95/15

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**

(71) Applicant: **BOEHRINGER INGELHEIM ESPANA
S.A.
Apartado 968
E-08080 Barcelona (ES)**

(72) Inventor: **Mato, José Maria
Jaén, 9
E-28230 Las Rozas (Madrid) (ES)
Inventor: Pajares, Maria Angeles**

**Infants Mercedes, 103
E-28020 Madrid (ES)**

Inventor: **Mingorance, Jesus
Gabriel y Galán, 2, 1 A
E-28320-Pinto (Madrid) (ES)**

Inventor: **Avarez, Luis
Islas Cies, 5 1D
ES-28035 Madrid (ES)**

(74) Representative: **Laudien, Dieter et al
Boehringer Ingelheim International GmbH
ZA Patente
Postfach 200
D-55216 Ingelheim am Rhein (DE)**

(54) **Production of S-adenosyl-methionine (SAM) by fermentation of transformed bacteria.**

(97) The present invention relates to a method of production of S-adenosyl-L-methionine (SAM) by fermentation of bacteria, especially bacteria transformed with an expression vector for SAM synthetase, expression vectors which can be used for said production, bacterial strains transformed with said vectors, and a method of producing said strains..




Fig. 2

EP 0 647 712 A1

The present invention relates to a method of production of S-adenosyl-L-methionine (SAM) by fermentation of bacteria, especially bacteria transformed with an expression vector for SAM synthetase, expression vectors which can be used for said production, bacterial strains transformed with said vectors, and a method of producing said strains.

SAM is a naturally occurring molecule, widely distributed throughout body tissues and fluids. It plays a central role in three main metabolic pathways, such as transmethylation, transsulphuration and aminopropylation. Through methylation, SAM regulates liver membrane lipid composition and fluidity, whereas by activating the transsulphuration pathway, it favors endogenous detoxifying processes.

The enzymatic machinery required for the synthesis of SAM is ubiquitous and occurs in almost all cells. SAM is synthesized from ATP and methionine, a reaction catalyzed by the enzyme S-adenosylmethionine synthetase (SAM synthetase, EC 2.5.1.6, Cantano, G. L. J. Biol. Chem. 1953, 204: 403-416).

SAM is currently used in medical and pharmacological areas due to its therapeutic potential in liver damage and affective disorders. A variety of clinical and experimental studies has demonstrated the ability of SAM to prevent or ameliorate the hepatotoxic effects of several drugs and chemicals (Friedel H. A. *et al.*, Drugs 1989, 38: 389-416; Corrales *et al.*, Drug Invest. 1992, 4, Suppl. 4: 8-13), to treat intrahepatic cholestasis of pregnancy (Frezza, M. *et al.*, Hepatogastroenterology 1990, 37: 122-125) and to improve clinical symptoms in cholestatic patients with chronic liver disease (Frezza, M. *et al.*, Hepatology 1987, 7: 1105; Frezza, M. *et al.*, Gastroenterology 1990, 99: 211-215). SAM has also been successfully used as an antidepressant (Baldessarini, R. J. *et al.*, Am. J. Med. 1987, 83 suppl. 5A: 95-103; Bell, K. M. *et al.*, Am. J. Psychiat. 1988, 145: 1110-1114; Chawla, R. K. *et al.*, Drugs 1990, 40 suppl. 3: 98-110). Increased therapeutic use of SAM has been made possible by the availability of a stable salt. However, the efficiency of the current methods of obtaining SAM is still low, despite attempts to produce it by enzymatic methods and/or conventional fermentation (Gross, A. *et al.*, Appl. Biochem. Biotechnol. 1983, 8: 415-422; Shiozaki, S. *et al.*, Agric. Biol. Chem. 1984, 48: 2293). Shiozaki *et al.* described the use of selected yeast strains which accumulated SAM for the production of the compound (Biotechnology & Bioengineering 1990, 35: 1120-1124).

The aim of the present invention is to provide an improved method of production of SAM as well as to provide means for performing this method. For this purpose, a bacterial strain suitable for producing high levels of SAM has been prepared. The process of preparation of this bacterial strain comprises the construction of a recombinant plasmid able to express rat liver SAM synthetase at high levels, its introduction into bacterial cells, growth of cells, induction of expression and isolation of SAM in a conventional manner.

The inducible prokaryotic expression vector used was pT7-7 (Tabor, S. & Richardson, C. C., Proc. Natl. Acad. Sci. USA 1985, 82: 1074-1078). This plasmid is available from Stan Tabor (Harvard Medical School, Boston, Ma). It contains a T7 RNA promoter upstream of the polycloning site, so that expression of the cloned fragment is directed by T7 RNA polymerase.

The bacterial strain used for transformation was *E. coli* BL21 (DE3) (F-ompT r^{-}_B m^{-}_B) (Studier, W. F. *et al.*, Methods Enzymol. 1990, 185: 60-89), which is available from William F. Studier (Brookhaven National Laboratory). This strain carries the gene for T7 RNA polymerase, which is expressed under the control of the *lacUV5* promoter. This promoter is inducible by isopropyl- β -D-thiogalactopyranoside (IPTG). Thus, the addition of IPTG induces the expression of T7 RNA polymerase, which, in turn, drives the expression of the cloned fragment in the plasmid pT7-7.

For constructing the recombinant plasmid, a 1.2 kb sequence of the cDNA clone pSSRL (Alvarez, L. *et al.*, FEBS Lett. 1991, 290: 142-146), corresponding to the coding region of rat liver SAM synthetase, was adapted for cloning into the *Nde*I/*Sal*I sites of the expression vector pT7-7 by means of the Polymerase Chain Reaction (PCR; Saiki, R. K. *et al.*, Science 1985, 230: 1350-1354). The resulting construct was designated pSSRL-T7N (Figure 1). The recombinant plasmid was introduced into *E. coli* BL21 (DE3) cells by the CaCl_2 method (Dagert, M. & Erlich, S. D., Gene 1974, 6: 23-28) and deposited under the accession number DSM 8592.

For monitoring SAM synthetase production, enzyme activity was assayed at different times after IPTG induction. Highest levels of SAM synthetase activity were obtained three hours after induction of the enzyme expression with IPTG (Figure 2). Consequently, the measurements of SAM levels were carried out three hours after the induction in either bacterial cells bearing pSSRL-TN7 or in the original strain. Non transformed *E. coli* BL21 (DE3) strain yielded an average of 0.1 nmol of SAM per ml of culture (1,2 nmol/mg protein), while the transformed strain yielded an average of 28 nmol of SAM per ml of culture (480 nmol/mg protein). Thus, a 300-fold increase of SAM content is observed in transformed cells, as compared with the original strain (Table 1). SAM can be isolated from the culture by known methods (Schlenk, F. *et al.* De Palma, R. E., J. Biol. Chem. 1957, 229: 1037-50; Cantoni, G. L., Methods Enzymol. 1957, 3: 600-3;

Schlenk, F. *et al.*, Arch. Biochem. Biophys. 1959, 83: 28-34; Svikla, G. *et al.* Schlenk, F., J. Bacteriol. 1960, 79: 841; Gordon, R. K. *et al.*, Methods Enzymol. 1987, 143: 191-5; Shiomi *et al.*, *loc. cit.*)

The method of the present invention has superior properties with respect to prior approaches to SAM production by fermentation of a microorganism. Bacterial strains obtainable as described in this specification therefore have a potential industrial interest as a powerful source of SAM.

Table 1

SAM content in transformed with pSSRL-T7N and nontransformed <i>E. coli</i> BL21 (DE3).		
Strain	nmol SAM/ml culture	nmol SAM/mg protein
<i>E. coli</i> BL21 (DE3)	0.12 ± 0.04	1.1 ± 0.5 (3) ^a
<i>E. coli</i> BL21 (DE3)/pSSRL-T7N	28.6 ± 6.4	482 ± 30 (3)

^aNumber of determinations

Figure legends

Fig. 1: Construction of the expression plasmid pSSRL-T7N.

A PCR fragment containing the coding sequence of rat liver SAM synthetase was digested by *Nde*I and *Xho*I restriction enzymes. The resulting fragment was then ligated to pT7-7 to obtain the recombinant plasmid pSSRL-T7N.

Fig. 2: SAM synthetase activity of *E. coli* BL21 (DE3) cells transformed with pSSRL-T7N, measured at different times after IPTG induction.

Examples

1. Construction of pSSRL-T7N

Two primers (5'-CGGAATCCATATGAATGGACCTGTGG 3') and (3'-GAACACAAAATC TCGGAGC-TCACG5') were synthesized with an Applied Biosystem 391 DNA synthesizer. The first one consisted of a sequence containing *Eco*RI and *Nde*I restriction sites at the 5'-end and a 16-nucleotide sequence at the 3'-end homologous to the bases 1-16 of the rat liver SAM synthetase clone pSSRL (Alvarez, L. *et al.*, FEBS Lett. 1991, 290: 142-146). The second one had a *Xho*I site at the 5'-end and a 15-nucleotide sequence complementary to the bases 1183-1197 of the pSSRL clone. The PCR was carried out in a reaction mixture containing 150 ng of pSSRL plasmid, 25 pmol each of the two oligonucleotides, 0.04 mM each of the four deoxyribonucleotides, 2 mM MgCl₂, 50 mM KCl, 10 mM Tris-HCl pH 9.0, 0.1% Triton 100X and 2.5 units of Taq DNA polymerase (Promega Inc., Madison, Wi) in a total volume of 50 µl of mineral oil (Sigma Chemical Co., St. Louis, Mo). Amplification was performed for 30 cycles at 94 °C for 2 min., 55 °C for 1 min., and 72 °C for 3 min, in a thermal cycler (Gene ATAQ controller, Pharmacia LKB). Thus, the amplified product contained the rat liver SAM synthetase coding region with a *Nde*I restriction site and *Xho*I site attached to the 5'- and 3'-ends of the cDNA, respectively. This product was then purified by phenol-chloroform extraction, digested with *Nde*I and *Xho*I and ligated to the *Nde*I-*Sal*I sites of the expression vector pT7-7, using T4 DNA ligase (Biotech SA, Madrid).

The ligation mixture was used to transform *E. coli* DH5α (relevant genotype: Δ*lac*U169, *hsd*R17, *rec*A1) using the method of CaCl₂ (Dagert, M. & Erlich, S. D., Gene 1974, 6: 23-28). Transformants were screened for plasmids containing the cDNA insert following digestion with *Nde*I and *Xho*I. One of these plasmids designated as pSSRL-T7N was then used to transform *E. coli* BL21 (DE3) cells. The resulting strain was deposited at the German Collection of Microorganisms and Cell Cultures (DSM), Mascheroder Weg 1b, D-38124 Braunschweig, Germany, under the accession number DSM 8592.

2. Culture of transformed BL21(DE3) and induction of SAM synthetase expression

An overnight culture was prepared from BL21(DE3) cells bearing the recombinant plasmid pSSRL-T7N. 1 ml of this culture was used to inoculate 100 ml LB medium (10 g bactotryptone, 5 g yeast extract, 10 g NaCl per liter) containing 100 ml/mg ampicillin. The cells were grown to A₅₉₅ 0.3-0.4 and isopropyl-β-D thiogalactopyranoside (IPTG) was added to a final concentration of 0.5 mM. Aliquots of 1 ml of the culture

were harvested by centrifugation at 1,2 and 3 hours after the initiation of induction. The pellets obtained were washed in water and resuspended in 100 μ l of 50 mM Tris pH 8.0, 1 mM EDTA, 10 mM $MgSO_4$ and protease inhibitors (2 μ g/ml aprotinin, 1 μ g/ml pepstatin A, 0.5 μ g/ml leupeptine, 2.5 μ g/ml antipain, 0.1 mM benzamidine, 0.1 mM PMSF) and sonicated. After centrifugation, the supernatants were used for measuring SAM synthetase activity.

3. Determination of SAM synthetase activity

SAM synthetase activity was assayed as described by Cabrero *et al.* (Eur. J. Biochem. 1987, 170: 299-304) using 160 μ l fraction samples and 90 μ l of a reaction mixture containing 75 mM Tris/HCl pH 8.0, 250 mM KCl, 9 mM $MgCl_2$, 10 mM dithiothreitol, 5 mM methionine and 5 mM $[2-^3H]ATP$ (4 Ci/mol). The incubation was carried out for 30 min at 37°C and was stopped by the addition of 3 ml distilled water. The solutions were immediately loaded on cation exchanger AG 50W-X4 columns (1 ml) equilibrated in water. The columns were washed with water (20 ml) and the $[^3H]SAM$ formed was then eluted with 4 ml 3 M ammonium hydroxide. The radioactivity was determined by counting in 10 ml of scintillation liquid in the presence of 1 ml glacial acetic acid.

Protein was determined by the method of Bradford (Bradford M. M., Anal. Biochem. 1977, 252: 248-254) using BSA as standard.

4. Measurement of SAM levels

SAM levels were determined in either bacterial cells bearing pSSRL-TN7 or non-transformed cells. Three hours after IPTG induction, aliquots of 1 ml of culture were immediately deproteinized by homogenization in 10% Trichloroacetic acid dissolved in 0.05 N HCl. TCA was eliminated by three successive washes with diethyl ether saturated with 0.05 N HCl, and samples were lyophilized. Samples were then resuspended in 0.01 M ammonium formate pH 4.0 and analyzed in a ultrasil CX HPLC column (4.6 mm x 25 cm; particle size 10 μ m) (Beckman, Palo Alto, Ca). After a 5 min washing period with 0.01 M ammonium formate pH 4.0, SAM was eluted in a 50 min gradient from 0.01 M to 0.8 M ammonium formate pH 4.0 at a flow rate of 1 ml/min. SAM was detected by measuring absorbance at 254 nm.

For measuring protein concentration, aliquots of 1 ml of the same cultures were centrifuged, resuspended in 100 μ l of 50 mM Tris pH 8.0, 1 mM EDTA, 10 mM $MgSO_4$, and sonicated. Protein was determined by the method of Bradford (Bradford M. M., Anal. Biochem. 1977, 252: 248-254) using BSA as standard.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT:
 (A) NAME: Boehringer Ingelheim Espana, S.A.
 (B) STREET: Apartado 14278
 (C) CITY: Barcelona
 (E) COUNTRY: Spain/Espana
 (F) POSTAL CODE (ZIP): 08080
 10 (G) TELEPHONE: (93)4045100
 (H) TELEFAX: (93)2042850
 (I) TELEX: 54722
- (ii) TITLE OF INVENTION: Production of S-Adenosyl-methionine (SAM) by
 fermentation of transformed bacteria
- 15 (iii) NUMBER OF SEQUENCES: 2
- (iv) COMPUTER READABLE FORM:
 (A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 20 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CGGAATCCAT ATGAATGGAC CTGTGG

26

35 (2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 40 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GCACTCGAGG CTCTAAAACA CAAG

24

50 Claims

1. Method of production of S-adenosyl-methionine by fermentation of a microbial host, characterized in that said host is a bacterium.
- 55 2. Method of claim 1, wherein said bacterium is overproducing S-adenosyl-methionine synthetase.
3. Method of claim 2, wherein said bacterium is transformed with an expression vector, said expression vector containing a nucleic acid sequence coding for S-adenosyl-methionine synthetase.

4. Method of claim 2, wherein said nucleic acid sequence is functionally linked to a bacteriophage T7 promotor.
5. Method of claims 1 to 4, wherein said bacterium is *Escherichia coli*.
6. Method of claim 5, wherein said *Escherichia coli* expresses bacteriophage T7 polymerase.
7. Method of claim 6, wherein said bacteriophage T7 polymerase expression is inducible.
8. Method of claims 2 to 7, wherein said S-adenosyl-methionine synthetase is from the rat.
9. Method of claims 5 to 8, wherein said *Escherichia coli* is the *Escherichia coli* strain deposited as DSM 8592.
10. Vector for expression of a gene in a bacterium, characterized in that it contains a nucleic acid sequence coding for S-adenosyl-methionine synthetase.
11. Vector of claim 10, wherein said nucleic acid sequence is functionally linked to a bacteriophage T7 promotor.
12. Vector of claims 10 to 11, which is a plasmid.
13. Vector of claims 10 to 12, wherein said S-adenosyl-methionine synthetase is from the rat.
14. Vector of claim 13, which is contained in the *Escherichia coli* strain deposited as DSM 8592.
15. Bacterial host, characterized in that it is transformed with a vector according to claims 10 to 14.
16. Bacterial host of claim 15, which is *Escherichia coli*.
17. Bacterial host of claim 16, wherein said *Escherichia coli* is the *Escherichia coli* strain deposited as DSM 8592.
18. Method of production of a bacterial strain suitable for production of S-adenosyl-methionine, characterized in that a bacterium is transformed with an expression vector, said expression vector containing a nucleic acid sequence coding for S-adenosyl-methionine synthetase.




Fig. 1




Fig. 2



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 11 6221

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.6)
X	EUROPEAN JOURNAL OF BIOCHEMISTRY vol. 184, no. 3, October 1989 pages 491 - 501 HORIKAWA SABURO ET AL. 'Isolation of a cDNA encoding the rat liver S-adenosylmethionine synthetase'	1-3,5,8, 10,13, 15,16,18	C12N15/54 C12P19/40 C12N15/70 C12N1/21 /(C12N1/21, C12R1:19)
Y	abstract * page 499, right column - page 500, left column *	4,6,7, 11,12	
D,X	FEBS LETTERS vol. 290, no. 1,2, September 1991, AMSTERDAM NL pages 142 - 146 LUIS ALVAREZ ET AL. 'Analysis of the 5' non-coding region of rat liver S-adenosylmethionine synthetase mRNA and comparison of the Mr deduced from the cDNA sequence and the purified enzyme'	1-3,5,8, 10,13, 15,16,18	
Y	* page 142, right column *	4,6,7, 11,12	TECHNICAL FIELDS SEARCHED (Int. CL.6) C12N C12P
D,Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA vol. 82, February 1985, WASHINGTON US pages 1074 - 1078 TABOR, S. ET AL. 'A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes' * page 1075, right column - page 1076, right column *	1-8, 10-13, 15,16,18	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16 March 1994	Examiner Espin, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 (11.82) (P04C01)

BEST AVAILABLE COPY



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 11 6221

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	JOURNAL OF BIOLOGICAL CHEMISTRY. (MICROFILMS) vol. 268, no. 19, 5 July 1993, BALTIMORE, MD US pages 13978 - 13986 SHIGEKI FUJIMOTO SAKATA ET AL. 'Cloning and expression of murine S-adenosylmethionine synthetase' * page 13979 - page 13982, left column; figure 2; table 1 *	1-8, 10-13, 15,16,18	
X	CHEMICAL ABSTRACTS, vol. 110, no. 19, 8 May 1989, Columbus, Ohio, US; abstract no. 171792d, FUJIO, TATSURO ET AL. 'Manufacture of S-adenosylmethionine with Brevibacterium, Corynebacterium, Escherichia, or Staphylococcus species' page 649 ; * abstract * & JP-A-63 279 798 (KYOWA HAKKO KOGYO CO., LTD.) 16 November 1988	1,2,5	
D,A	BIOTECHNOLOGY AND BIOENGINEERING. INCLUDING: SYMPOSIUM BIOTECHNOLOGY IN ENERGY PRODUCTION AND CONSERVATION vol. 35, 1990, NEW YORK US pages 1120 - 1124 NAOFUMI SHIOMI ET AL. 'Production of S-Adenosyl-L-Methionine by Saccharomyces cerevisiae cells carrying a gene for ethionine resistance' * the whole document *		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16 March 1994	Examiner Espen, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	

EPO FORM 1503 (11.82) (P04C01)

BEST AVAILABLE COPY



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 11 6221

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,A	AGRIC. BIOL. CHEM. vol. 48, no. 9, 1984 pages 2293 - 2300 SHOZO SHIOZAKI ET AL. 'Unusual intracellular accumulation of S-adenosyl-L-methionine by microorganisms' * the whole document * -----		
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16 March 1994	Examiner Espen, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1501 (01.92) (P0400)

BEST AVAILABLE COPY